

## Impact of climate change on the western Himalayan mountain ecosystems: An overview

G. C. S. NEGI<sup>1\*</sup>, P. K. SAMAL<sup>2\*</sup>, J. C. KUNIYAL<sup>3</sup>, B. P. KOTHYARI<sup>1</sup>, R. K. SHARMA<sup>3</sup> & P. P. DHYANI<sup>1</sup>

<sup>1</sup>*G. B. Pant Institute of Himalayan Environment & Development (GBPIHED)  
Kosi-Katarmal, Almora 263 643, Uttarakhand*

<sup>2</sup>*GBPIHED, North East Unit, Itanagar 791 113, Arunachal Pradesh*

<sup>3</sup>*GBPIHED, Himachal Unit, Kullu 175 126, Himachal Pradesh*

**Abstract:** This article presents an overview of climate change impacts on agriculture, water and forest ecosystems in the western Himalayan mountains based on literature review and some anecdotal evidences. A great deal of research work has been carried out on different aspects of western Himalayan mountain ecosystems but the findings have yet to be correlated in the context of climate change. There is a need to strengthen climate data collection network which is presently insufficient to meet the requirement of climate change research. The climate data in the region is scarce and in many instances does not involve uniform methodology and standard instrumentation. The data reliability thus is uncertain as the data are based on crude collection methods without quality control. Climate change impacts also need to be categorized according to various climatic elements viz., rainfall, temperature, CO<sub>2</sub> concentration, etc. Coordinated efforts are required for adaptation and mitigation as the vulnerable mountain ecosystems and communities are likely to face greater risk of climate change impacts than other ecosystems. Research and documentation is also required to validate the indigenous methods of adaptations and coping up mechanisms. Capacities of communities have to be enhanced and strategies are to be developed for adaptation to climate change. There is also a need to network with other potential players in this subject to utilize the synergy in the best interest of survival, and ensuring livelihood security of the inhabitants of the region and that of the adjacent lowlands. The balance between economic interests and ecological imperatives is also essential.

**Resumen:** Este artículo presenta una visión integral de los impactos del cambio climático sobre la agricultura, el agua y los ecosistemas forestales en los Himalaya Occidentales, con base en una revisión de literatura y algunas evidencias anecdóticas. Se ha hecho mucha investigación sobre diferentes aspectos de los ecosistemas de montaña de los Himalaya Occidentales, pero los resultados no se han correlacionado en el contexto del cambio climático. Es necesario fortalecer la red de recolección de datos de clima, ya que actualmente ésta es insuficiente para cubrir los requerimientos de la investigación sobre el tema. Los datos de clima en la región son escasos y con frecuencia no involucran el uso de instrumentación estándar y metodología uniforme. La confiabilidad de los datos es incierta, pues se desprenden de un conjunto burdo de métodos carentes de control de calidad. Asimismo, es necesario categorizar los impactos del cambio climático de acuerdo con varios elementos climáticos, p.ej. precipitación pluvial, temperatura, concentración de CO<sub>2</sub>, etc. Hacen falta esfuerzos coordinados para la adaptación y la mitigación porque es probable que los vulnerables ecosistemas de montaña enfrenten mayores riesgos de ser impactados por el cambio climático que otros ecosistemas. También se necesita investigación y documentación para validar los métodos nativos de adaptación y los mecanismos para hacer frente al cambio, así como incrementar las capacidades

---

\* *Corresponding Author*; e-mail: gcsnegi@yahoo.co.in

de las comunidades y desarrollar estrategias de adaptación. También hay que establecer redes con otros actores potenciales en este temapara utilizar la sinergia en el mejor interés por la supervivencia, y asegurar la seguridad de la subsistencia de los habitantes de la región y de las tierras bajas adyacentes. También es esencial conseguir un balance entre los intereses económicos y las prioridades ecológicos.

**Resumo:** Com base numa revisão da literatura e algumas evidências anedóticas,este artigo apresenta uma visão geral dos impactes das mudanças climáticas na agricultura, água e nos ecossistemas florestais no oeste das montanhas dos Himalaias. Uma grande parte do trabalho de pesquisa foi realizado em diferentes aspectos dos ecossistemas nas montanhas ocidentais dos Himalaias, se bem que os resultados obtidos ainda não tenham sido correlacionados no contexto das mudanças climáticas. Há uma necessidade de reforçar a rede de coleta de dados climáticos, que é atualmente insuficiente, para atender às exigências de pesquisa sobre as mudanças climáticas. Os dados climáticos na região são escassos e, em muitos casos, não são gerados por metodologias uniformes e instrumentação estandardizada. A fiabilidade dos dados é assim incertajá que os dados têm por basemétodos grosseiros de recolha sem controle de qualidade. Os impactes das mudanças climáticas também necessitam ser categorizadosde acordo com vários elementos climáticos i.e. precipitação, temperatura, concentração de CO<sub>2</sub>, etc.. São ainda necessários esforços coordenados para adaptação e mitigação já que a vulnerabilidade dos ecossistemas de montanha e as comunidades são provavelmente as que enfrentam maior risco dos impactos das mudanças climáticas em confronto com outros ecossistemas. A pesquisa e documentação são também necessárias para validar os métodos indígenas de adaptação e mecanismos de como os enfrentam. As capacidades das comunidades têm que ser reforçadas e as estratégias para a adaptação às mudanças climáticas devem ser desenvolvidas. Neste assunto há também uma necessidade de interagir com outros interventorespotenciais por forma a utilizar sinergias no melhor interesse da sobrevivência, subsistência e assegurar a segurança do modo de vida dos habitantes da região e das terras baixas adjacentes. O equilíbrio entre interesses económicos e os imperativos ecológicos é também essencial.

**Key words:** Adaptation and mitigation, climate change, Himalayan ecosystems, impacts, preparedness, R & D needs, vulnerability.

## Introduction

Climate change (CC) is a major challenge facing our planet today. This is an all-encompassing threat that will pose significant environmental, economic, social and political challenges for years and decades to come. Climate change has emerged as a global environmental issue that has engaged the world attention as it relates to global common atmosphere. It is scientifically least predictable, and its impacts are likely to affect adversely the vulnerable and poor people mostly, who have contributed least to the major causes of CC. Mountains are early indicators of climate change (Singh *et al.* 2010). As glaciers recede, and snowlines move upwards, river flows are likely to change, and alteration in water flow regime may lead to a plethora of social issues and affect hydro-

power generation, endanger biodiversity, forestry and agriculture-based livelihoods and overall well-being of the people.

### *Climate change impacts on the Himalayan ecosystems*

At the outset it may be pointed out that research on CC vis-à-vis its impact on ecosystems (e.g., forests, water, agricultural resources, etc.) is still in an infancy stage in the Himalayan mountains. As such there is a paucity of long-term climate data in the region, and data recorded do not employ uniform instrumentation and methodology hence the data quality is uncertain (Joshi & Negi 1995). Very limited studies are available which could demonstrate the impact of CC on the mountain ecosystems. However, with the release of recent IPCC report research on CC is occupying

the agenda of many institutions. Preliminary studies indicate that the Himalaya seems to be warming more than the global average rate (Liu & Chen 2000; Shrestha *et al.* 1999), temperature increases are greater during the winter and autumn than during the summer; and the increases are larger at higher altitudes (Liu & Chen 2000). The Indian Institute of Tropical Meteorology, Pune has reported a decrease in precipitation over 68 per cent of India's area over the last century (Kumar *et al.* 2006). However, significant increase in rainfall was noticed in Jammu and Kashmir and some parts of Indian peninsula (Agarwal 2009). A study indicates that the mean annual temperature in the Alaknanda valley (western Himalaya) has increased by 0.15 °C between 1960 - 2000 (Kumar *et al.* 2008a). Satellite imagery suggests almost 67 % of the glaciers in the Himalaya have retreated (Ageta & Kadota 1992). For example, in Nepal this process is as fast as 10 m year<sup>-1</sup> (Ageta & Kadota 1992). The average temperature of Kashmir valley has gone up by 1.45 °C over the last two decades (Sinha 2007). Impact of changing climate is also perceptible on vegetation. In some parts of the high altitude, the biodiversity is being lost or endangered because of land degradation and the over use of resources, e.g., in 1995, about 10 % of the known species in the Himalaya were listed as 'threatened' (IPCC 2002). However, impact of CC on biodiversity and vegetation in the region is yet to be carefully studied to establish such a relationship.

Exponential increase in green house gases (GHGs) like carbon dioxide, methane, nitrous oxide, water vapour, CFCs, etc., in the atmosphere has resulted in CC change (World Climate News 2006). The concentration of CO<sub>2</sub>, mainly responsible for global warming, has reached to 379 ppm in 2005 from its pre-industrial value (i.e., 280 ppm). The increase in GHGs between 1970 and 2004 was approximately 70 %. The mean temperature of the earth has increased by 0.74 °C during last century (World Climate News 2006). Globally, the sea level rose at the rate of 1.8 mm year<sup>-1</sup> during 1961 to 2003, and faster (i.e., at the rate of 3.1 mm year<sup>-1</sup>) during 1993 to 2003. The review report (Pasupalak 2009) projecting the scenarios of global warming indicate that the global average surface temperature could rise by 1.4 to 5.8 °C by 2100. Global mean sea level is projected to rise by 0.18 to 0.59 m by the end of the current century.

This article provides a brief overview of impacts on different components (people, agricul-

ture, water and forest ecosystems) in the western Himalayan mountains based on literature review and some anecdotal evidences. Some important research areas for further investigation are also listed.

### *Socio-economic and health impacts*

Climate change can influence the socio-economic setting in the Himalaya in a number of ways. It can influence the economy (e.g., agriculture, livestock, forestry, tourism, fishery, etc.) as well as human health. Specific knowledge and data on human wellbeing in the Himalaya is limited, but it is clear that the effects of CC will be felt by people in their livelihoods, health, and natural resource security, among other things (Sharma *et al.* 2009). The consequences of biodiversity loss from CC are likely to be the worst for the poor and marginalized people who depend almost exclusively on natural resources. Poverty, poor infrastructure (roads, electricity, water supply, education and health care services, communication, and irrigation), reliance on subsistence farming and forest products for livelihoods, substandard health indicators (high infant mortality rate and low life expectancy), and other indicators of development make the Himalaya more vulnerable to CC as the capacity to adapt is inadequate among the inhabitants.

Climate change has a powerful impact on human health directly (e.g., impacts of thermal stress, death/injury in floods and storms) and indirectly through changes in the ranges of disease vectors, water-borne pathogens, water quality, air quality, food availability and quality (McMichael *et al.* 2003), cardiovascular mortality and respiratory illnesses, transmission of infectious diseases and malnutrition from crop failures (Patz *et al.* 2005). Climatic variations are likely to have a direct impact on the epidemiology of many vector-borne diseases. It is projected that the spread of malaria, bartonellosis, tick-borne diseases and other infectious diseases linked to the rate of pathogen replication will all be enhanced. Several studies using statistical models based on empirical weather data have concluded that CC impacts epidemics of vector-borne disease, such as Ross Virus Fever in Australia (Woodruff & Guest 2002). Climate change and variability are likely to highly influence current vector-borne diseases epidemiology as it is estimated that by the year 2100, average global temperatures will have increased by 1.0 - 3.5 °C (Watson *et al.* 1997), increasing the

likelihood of many vector-borne diseases. With a rise in surface temperature and changes in rainfall patterns, the distribution of vector mosquito species may change (Patz & Martens 1996; Reiter 1998). Malaria mosquitoes have recently been observed at high altitudes in the region (Eriksson *et al.* 2008). Temperature can directly influence the breeding of malaria protozoa and suitable climatic conditions can intensify the invasiveness of mosquitoes (Tong & Ying 2000). Another concern is that changes in climate may allow more virulent strains of disease or more efficient vectors to emerge or be introduced to new areas (Sharma *et al.* 2009).

Aerosols have been considered the primary cause for the increase of the Earth's atmospheric temperature. In Himachal Pradesh, aerosols optical depth (AOD), obtained through Multi-wavelength Radiometer (MWR) has shown highest ever AOD at 500 nm as  $0.55 \pm 0.03$  in May 2009 which was 104 % more than mean AOD value from April 2006 to December 2009 (Kuniyal *et al.* 2009). This value of AOD was found to be  $0.056 \pm 0.037$  at Nainital (Pant *et al.* 2006), that is, far less to the values obtained at Kullu, indicating inter-regional variations in CC within the Himalayan region. Temperature rise due to radiative forcing from aerosols in the atmosphere based on per unit AOD increase at Mohal-Kullu (HP) was calculated as high as 0.95 kelvin (K) day<sup>-1</sup> during summer (April-July) and as low as 0.51 K day<sup>-1</sup> during winter season (December, January-March) (Guleria *et al.* 2010).

Climate change will have a wide range of health impacts across the Himalaya. For example, increase in malnutrition due to the failure of food supply, disease and injury due to extreme weather events (Epstein *et al.* 1995), increase in diarrhoea diseases from deteriorating water quality, increase in infectious diseases and cardio-respiratory diseases from the build-up of high concentrations of air pollutants such as nitrogen dioxide (NO<sub>2</sub>), lower tropospheric and ground-level ozone, and air-borne particles in large urban areas. Huge quantities of municipal waste produced in the western Himalayan mountain towns are further compounding the problem of emission, sanitation and associated health hazards (Kuniyal 2002).

#### *Impact on agro-ecosystems*

Agriculture is highly dependent on weather, and changes in weather cycle have a major effect on crop yield and food supply. Mountain agriculture is mostly rainfed (appx. 85 %), and driven

by biomass energy of surrounding forests and confined to terraces carved out of hill slopes. The small holdings (< 1 ha in majority of cases) are distributed over small parcels of land and the agricultural productivity is very low (6 - 13 q ha<sup>-1</sup>). The irrigation systems, have been severely affected with the rainfall becoming erratic. In the Kullu valley (HP) it has been reported that the rainfall has decreased by about 7 cm, snowfall by about 12 cm, but the mean minimum and maximum temperatures have increased by 0.25 - 1 °C, respectively in 1990s as compared to 1880s (Vishvakarma *et al.* 2003). Various studies have reported that the rate of apple production in Kullu valley has significantly declined during the period 1981-2000 (Vishvakarma *et al.* 2003; Vedwan & Rhoades 2001). These workers have observed that apple cultivation has shifted to higher altitudes and apple yield mainly in lower altitudes has declined due to inadequate chilling as the temperature at lower altitudes is rising due to CC in Himachal Pradesh. Because of the change in snowfall the chilling hours for apple trees are reduced, affecting the time of bud-break. Early snow (December to early January) is preferred for its favourable effect on bud-break and soil moisture. It provides a chilling period of about 10 weeks below 5 °C, which is required to meet the internal condition necessary for bud-break in apples in springtime (Abbott 1984).

Some of the documented impacts on mountain agriculture that are linked with CC in the Himalayan region are: (i) Reduced availability of water for irrigation; (ii) Extreme drought events and shifts in the rainfall regime resulting into failure of crop germination and fruit set; (iii) Invasion of weeds in the croplands and those are regularly weeded out by the farmers (e.g., *Lantana camara*, *Parthenium odoratum*, *Eupatorium hysterophorus* etc.); (iv) Increased frequency of insect-pest attacks; (v) Decline in crop yield (Negi & Palni 2010). These factors have led to loss in agri-diversity and change in crops and cropping patterns. Traditional agriculture in the Himalayan mountains has been a rich repository of agrobiodiversity and resilient to crop diseases. For example, in Uttarakhand over 40 different crops and hundreds of cultivars selected by farmers, comprising cereals, millets, pseudo-cereals, pulses and tuber crops are cultivated (Agnihotri & Palni 2007; Maikhuri *et al.* 1997). Mixed cropping of 12 crops (*Baranaja*) is another best example of rich agri-diversity of the region (Ghosh & Dhyani 2004). These crops are adapted to the local environmental

conditions and possess the inherent qualities to withstand the environmental risks and other natural hazards. This adaptability has ensured the food and nutritional security of the hill farmers from generations. However, the area under traditional crops has drastically declined (> 60 %) particularly during the last three decades and many of the crops are at the brink of extinction, such as *Glycine* spp., *Hibiscus sabdariffa*, *Panicum miliaceum*, *Perilla frutescens*, *Setaria italica*, *Vigna* spp., to name a few (Maikhuri *et al.* 2001; Negi & Joshi 2002). Recent studies at the Indian Agriculture Research Institute (IARI), New Delhi indicate the possible loss of 4 - 5 million tons in wheat production in future with every rise of 1 °C temperature throughout the growth periods. Losses in other crops are still uncertain but they are expected to be relatively smaller, especially for kharif crops (Uprety & Reddy 2008). Deficit in food production in Kashmir region has reached 40 % in 2007 from 23 % in 1980 - 81 has been linked with CC (Sinha 2007).

Alterations in the floral diversity due to landuse and land cover change and extinction of local cultivars will also affect the population of pollinators. In the Himalaya honey bees have been the important component of mountain agriculture as they provide honey with nutrition and medicinal value. Recent reports reveal that due to habitat loss through landuse changes, increasing monoculture and negative impacts of pesticides and herbicides, the diversity of native bees is declining in the region (Pratap 1999). Studies carried out by International Centre for Integrated Mountain Development (ICIMOD) reveal that declining apple productivity is a result of inadequate pollination (Pratap & Pratap 2003). The farmers are now compelled to rent colonies of honey bees for pollinating the apple orchards @ Rs. 500/colony (Ahmad *et al.* 2002), and devise short-term solutions until the “polliniser” trees in their orchards begin flowering. Bunches of small flowering branches of the pollinisers called “bouquets” are put in plastic bags filled with water and hung on the branches of commercially premium apple varieties to attract the pollinators.

Climate change also leads to shift in pest incidence, migration and viability. A change in climatic conditions can cause a pest or disease to expand its normal range into a new environment, extending losses and affecting natural plant communities (Rosenzweig *et al.* 2001). Temperature is one of the dominant factors affecting the growth rate and development of insect pests (Bale

*et al.* 2002; Patterson *et al.* 1999). High summer temperatures would favour growth of temperate zone insects leading to faster development and additional generations per year (Bale *et al.* 2002; Porter *et al.* 1991). High winter temperatures are likely to result into increased over-wintering thereby increasing population levels in the following season (Porter *et al.* 1991). It was found that aggressiveness (defined as the percentage of leaf area of a diseased plant) of the fungal pathogen *Colletotrichum gloeosporioides* increased under CO<sub>2</sub> concentration double than the ambient level (Chakraborty & Datta 2003). More extreme weather events such as droughts and floods are likely to lead to pest and disease outbreaks (Rosenzweig *et al.* 2001). Drought stress affects plant physiology, causing some plants to become more susceptible to pests and pathogens, especially when combined with higher temperatures, which can suppress plant defense responses (Rosenzweig *et al.* 2001).

Many of the crop species in the Himalaya vary in their response to CO<sub>2</sub>. C<sub>3</sub> crops such as wheat, rice, and soybeans respond readily to increased CO<sub>2</sub> levels. Corn, sorghum, sugar-cane, and millet are C<sub>4</sub> plants that follow a different pathway. The initial short-term studies indicated that photosynthesis is stimulated more in C<sub>3</sub> species as compared to C<sub>4</sub> species in response to CO<sub>2</sub> enrichment (Uprety & Reddy 2008). A preliminary study (Joshi & Palni 2005) relating to response of crops to elevated CO<sub>2</sub> and temperature in the Himalayan mountains shows that *Hordeum himalayans* responds negatively to elevated atmospheric temperature and reduces photosynthesis by 25 - 30 % and consequently crop yields are low. However, the other species, *H. vulgare*, shows a reverse trend. Thus far, these effects have been demonstrated mainly in controlled environments such as growth chambers, greenhouses, and polyhouses. Increased evaporation from the soil and accelerated transpiration in the plants themselves will cause moisture stress; as a result there will be a need to develop crop varieties with greater drought tolerance (Rosenzweig & Hillel 1995). The positive effects of CC – such as longer growing seasons and faster growth rates at higher altitudes – may be offset by negative factors such as changes in established reproductive patterns, migration routes and ecosystem (Sharma *et al.* 2009).

#### *Impact on forest ecosystems*

Vegetation patterns (distribution, structure and ecology of forests) across globe are controlled

mainly by the climate. Several climate-vegetation studies have shown that certain climatic regimes are associated with particular plant communities (Thornthwaite 1948; Walter 1985; Whittaker 1975). The Third Assessment Report of IPCC (2001) concluded that the forest ecosystems could be seriously impacted by future CC. Even with global warming of 1 - 2 °C, much less than the most recent projections of warming during this century (IPCC 2001a), most ecosystems and landscapes will be impacted through changes in species composition, productivity and biodiversity (Leemans & Eickhout 2004). It has been demonstrated that upward movement of plants will take place in the warming world (Cannone *et al.* 2007; Kelly & Goulden 2008; Pauli *et al.* 2003). Due to increase in temperatures, change in vegetation, rapid deforestation and scarcity of drinking water, habitat destruction and corridor fragmentation may lead to a great threat to extinction of wild flora and fauna.

In the western Himalayan mountains early flowering of several members of Rosaceae (e.g., *Pyrus*, *Prunus* spp.) and Rhododendrons has often been linked with global warming. Negi (1989) found that forest trees along an altitudinal gradient of 600 - 2200 m altitude in Kumaun Himalaya vary with respect to periodicity of phenophases such as vegetative bud break, flowering, fruiting and leaf drop. These phenophases were found to be influenced by variations in temperature and rainfall changes over two consecutive years (1985-1987). For example, at approx. 2000 m altitude leaf initiation was advanced by a week in 1985 (spring temperature in March and April were 17.6 & 20.9 °C, respectively) as compared to that in 1986 (spring temperature in March = 14.8 °C and April = 16.5 °C). However, this annual shift in occurrence of phenophases was not observed for all the species. In the year 1985 when the spring temperature was higher as compared to 1986 flowering was also advanced by 1-3 weeks (in tree species like *Quercus* spp., *Rhododendron arboreum*, *Pinus roxburghii* - the dominant forest trees) in 1985 as compared to 1986 (Negi 1989). It appears that plant taxa are impacted differently by CC calling for further detailed investigations.

Spread of alien invasive species such as *Lantana*, *Eupatorium* and *Parthenium* spp.) in the natural forests has also been linked with CC, which will have a competitive impact on existing species. Research on alpine vegetation suggests that many species are able to start their growth with the supply of snow-melt water well before the monsoon begins in June (Negi *et al.* 1991). Growth

and life cycles of these species are already being disturbed because of reduced water from snow melt.

Increased incidences of forest fire are another prominent change that is linked with CC. The frequency, size, intensity, seasonality, and type of fires depend on weather and climate in addition to forest structure and composition. In 2005-2006, a record 8,195 hectares of forest in Himachal Pradesh was lost to fire (Bhatta 2007). In four districts of Uttarakhand (Almora, Chamoli, Tehri and Pauri Garhwal), in one of the most devastating forest fires of recent decades on 27 May, 1995 a total of 2115 km<sup>2</sup> (between altitudes 600 m to 2650 m) was damaged severely (Semwal & Mehta 1996). A comparison of air temperatures during the fire season for the two years 1994 (no fire event) and 1995 (large fire events) in Binsar Wildlife Sanctuary, Kumaun Himalaya showed that in 1995 the air temperature was higher (Sharma & Rikhari 1997). It is expected that with the CC, scenario of the forests, both in terms of structure and functioning, is likely to change substantially. In the case of many dominant forest species of the region like sal (*Shorea robusta*), tilonj oak (*Quercus floribunda*) and kharsu oak (*Q. semecarpifolia*) seed maturation and seed germination coincide with monsoon rainfall. In wet conditions these species show vivipary. A rise in temperature and water stress may advance seed maturation, which might result in the breakdown of synchrony between monsoon rains and seed germination (Singh *et al.* 2010).

Forests are also the source of NTFPs (including medicinal plants). The oak forests of the region are the storehouse of biodiversity, NTFPs and a variety of medicinal plants. For example, *Morchella esculenta* (an edible fungi) that is found in the oak-conifer forests of this region in the Niti valley of Chamoli distt. is collected by the local people. A family on an average collects about 1.5 kg dry wt. of *Morchella* every year which is sold @ Rs. 5000/kg (Prasad *et al.* 2002). Another such example is that of Kafal (*Myrica esculenta*) fruits collected by the local people from the forests that are sold raw in the nearby markets @ Rs. 20-40/kg. It has been estimated that in the Kumaun region alone people can earn appx. Rs. 1.4 million per season from selling this fruit (Bhatt *et al.* 2000). It is expected that with the increase in drought cycles and concomitant increase in forest fires the pine (*Pinus roxburghii*) forests will encroach upon the area under oak (*Quercus* spp.) forests and also reduce the yield of non-timber forest products

(NTFPs). Apart from the direct benefits accrued from these forests the ecosystem services generated would also alter (Semwal *et al.* 2007). Inevitably, any change in the forest (distribution, density and species composition) under CC would immensely influence economies like forestry, agriculture, livestock husbandry, NTFPs and medicinal plants based livelihoods and many others.

World over, many studies have been conducted on the effects of elevated CO<sub>2</sub> on photosynthesis and yield of plants. The so called "CO<sub>2</sub> fertilization effect" is often examined using greenhouses, growth chambers and open top chambers (Nowak *et al.* 2004; Pendall *et al.* 2004). In the western Himalaya, oak (*Quercus* spp.) and other fodder trees (e.g., *Grewia*, *Celtis*, *Boehmeria* spp.) show over 50 % higher photosynthesis under elevated CO<sub>2</sub>, whereas *Betula utilis* (a tree of alpine habitat) exhibited a reverse trend (Joshi *et al.* 2007). In a recent study (Ravindranath *et al.* 2008) it was reported that under two scenarios of 740 ppm CO<sub>2</sub> (A<sub>2</sub>) and 575 ppm CO<sub>2</sub> (B<sub>2</sub>), vegetation response model for the forests of India projected for the year 2085, showed that 77 % and 68 % of the forested grids are likely to experience shift in forest types under A<sub>2</sub> and B<sub>2</sub> scenarios, respectively. The impacts of CC on forest ecosystems include shifts in the latitude of forest boundaries and the upward movements of tree line to higher elevations; changes in species composition and in vegetation types; and an increase in net primary productivity (Ramakrishna *et al.* 2003).

Increase in temperature is known to enhance a ground-level ozone concentration which is one of the GHGs and a phyto-toxic air pollutant that leads to serious injury in plant tissues and reductions in their growth and productivity (USEPA 1996). It has been shown that exposure to O<sub>3</sub> concentration of more than 50 ppb is sufficient to cause damage to certain species (Aneja *et al.* 1991). Ozone exposure to plants during the day brings about biochemical changes resulting in decreased photosynthesis (Musselman & Massman 1999). In Himachal Pradesh, ozone episodes have shown 86.7 ppb at 1500 h IST at Mohal (Kuniyal *et al.* 2007), which increased up to 95.19 ppb at 1700 h in 2007.

#### *Impact on water resources*

The Himalayan region represents one of the most dynamic and complex mountain systems and extremely vulnerable to global warming (Bandyopadhyay & Gyawali 1994). The Himalayan mountains

are important sources of water to the Indo-Gangetic plains through the perennial glacier fed rivers. During the Pleistocene era (2 million years ago) glaciers occupied about 30 % of the total area of the Earth as against 10 % at present (Bahadur 1998). The IPCC has stated that thinning of glaciers since the mid-19th century has been obvious and pervasive in many parts of the world. The reduction in ice cover during the last century, across the globe, especially in mountain glaciers is seen as evidence of CC (IPCC 2001 b). At high elevations in the Himalaya, an increase in temperature could result in faster recession of glaciers and an increase in the number and extent of glacial lakes - many of which have formed in the past several decades. The rapid growth of such lakes could exacerbate the danger from glacial lake outburst floods (GLOFs), with potentially disastrous effects.

Studies on selected glaciers of Indian Himalaya indicate that most of the glaciers are retreating discontinuously since post-glacial time. The Gangotri glacier, one of the major and important glaciers in the Himalaya, was 25 km long when measured in 1930s and has now shrunk to about 20 km (Hasnain 1999). The average rate of recession of this glacier between 1985 and 2001 is about 23 m per year (Hasnain 2002). Thakur *et al.* (1991) have reported the retreat of Gangotri glacier at an average rate of 18 m per year. While with the use of Differential Global Positioning System, it has been reported that the Gangotri glacier has retreated at much lower rates (11.80 ± 0.035 m per year) between the years 2005 - 2007 (Kumar *et al.* 2008b; Raina 2009). The Siachen & Pindari glaciers retreated at the rates of 31.5 m and 23.5 m per year, respectively (Vohra 1981). Shukla & Siddiqui (1999) monitored the Milam Glacier in the Kumaon Himalaya and estimated that the ice retreated at an average rate of 9.1 m per year between 1901 and 1997. Dobhal *et al.* (1999) monitored the shifting of snout of Dokriani Bamak Glacier in the Garhwal Himalaya and found 17.2 m per year retreat during the period 1962 to 1997. The average retreat was 16.5 m per year. Matny (2000) found Dokriani Bamak Glacier retreated by 20 m in 1998, compared to an average retreat of 16.5 m over the previous 35 years. Geological Survey of India studied the Gara, Gor Garang, Shaune Garang, Nagpo Tokpo Glaciers of Satluj River Basin and observed an average retreat of 4.22 - 6.8 m year<sup>-1</sup> (Vohra 1981). The Bara Shigri, Chhota Shigri, Miyar, Hamtah, Nagpo Tokpo, Triloknath and Sonapani Glaciers in

Chenab River Basin retreated at the rate of 6.81 to 29.78 m year<sup>-1</sup>. A massive glacial retreat rates of 178 m year<sup>-1</sup> in Parbati Glacier in Kullu District during 1962 to 2000 has been observed (Kulkarni *et al.* 2004). These observations, irrespective of the differences in the retreat of glaciers, suggest that global warming and CC has affected snow-glacier melt and runoff pattern in the Himalaya.

When assessing the warming effects on glacier melting in the Himalaya, the role of black carbon (BC) cannot be neglected (Ming *et al.* 2008). In the Himalayan region, solar heating from BC at high elevations may be just as important as carbon dioxide in the melting of snowpacks and glaciers (Ramanathan & Carmichael 2008). The first historical record of BC deposition of about 80 ng m<sup>-3</sup> during 1951 - 2001 in a 40-m shallow ice core retrieved from the East Rongbuk Glacier in the northeast saddle of Mt. Qomolangma (Everest) during the past ~50 yrs in the high Himalaya (Ramanathan & Carmichael 2008). The rise in BC in South Asia could penetrate into the Tibetan Plateau by climbing over the elevated Himalaya. In the northwestern Indian Himalayan region, BC concentration at Mohal-Kullu (H.P.) showed bimodal peaks- one at 0600 to 0900 h IST and another at 1800 h IST showing hourly average value always above 2500 ng m<sup>-3</sup> (Kuniyal 2010). However, highest ever value on hourly basis was 15657 ng m<sup>-3</sup> in January 2010 followed by 15006 ng m<sup>-3</sup> in December 2009. In the morning hours due to shallow boundary layer in the hills the BC recorded a peak. Again in the evening the BC recorded a peak value due to dispersion of C produced by biomass burning, vehicular emission and other aerosols (Kuniyal 2010).

Mountain springs have been reported to decline the water yield or have gone dry mainly due to the erratic rainfall in the recent decades (Valdiya & Bartarya 1989 a; Negi & Joshi 2004). A survey in the Gaula river catchment in the Central Himalaya revealed that about 45 % of springs have gone dry or become seasonal (Valdiya & Bartarya 1989 a). Studies on aquifer response to regional climate variability in a part of Kashmir Himalaya (Jeelani 2008) revealed that out of forty major perennial springs monitored under different lithological controls, the average monthly spring discharge is high in Triassic Limestone-controlled springs (Karst springs) and low in alluvium- and Karewa-controlled springs. In general, the measured monthly spring discharges show an inverse relation with the monthly precipitation data. However, a direct correlation exists between the

spring discharges and the degree of snow/ice melt. Rainfall patterns have been changed in the recent decades. The monsoon rains are delayed and continuing rainfall for seven days (*Satjhar*)- a phenomenon which was a characteristic feature of monsoon season has almost disappeared now. Similarly, the winter rains have also become unpredictable and lower in quantity. In many areas, a greater proportion of total precipitation appears to be falling as rain than before. As a result, snowmelt begins earlier; this affects river regimes, drought and flood cycles, natural hazards, water supplies, and people's livelihoods and infrastructure. Erosion has increased five-fold to 1 mm per year (Valdiya & Bartarya 1989b). The rivers carry large amounts of sediment at rates of 16.5 ha-meter per 100 km<sup>2</sup> of the catchment area per year (Valdiya & Bartarya 1989b).

#### *Research & development issues for future action*

Considering the potential impacts of CC on the Himalayan ecosystem following few important areas need attention both of the researchers and policy planners. Some of the important areas of research could be: (i) Collection of meteorological data involving standard methodology and instrumentation, (ii) Understanding drought and flood cycles, climate variability and other extreme events, (iii) Plant taxa vulnerable to CC and survival strategies of plants, (iii) Global warming associated with upward migration of altitudinal boundaries and consequent change in snowline position and its biota, (iv) Habitat requirements and corridors for upward migration of plants and animal species, (v) Impact of CC (involving eco-physiological studies such as drought, elevated CO<sub>2</sub> and atmospheric temperature) on important food crops, timber spp., medicinal plants which may affect the food security and revenue generation for the region, (vi) Studies on air pollutants (AODs, GHGs) and temperature rise to observe the impact of aerosols, GHGs and other pollutants on the atmospheric temperature, (vii) Invasion of weeds, such as *Lantana*, that compete with the native flora for soil nutrients and (viii) Documentation of local traditional knowledge of climate variability and coping-up strategies can be useful in formulating strategies to adapt to the impact of CC.

#### **Acknowledgements**

Authors are thankful to Dr. L.M.S. Palni, Director of the G. B. Pant Institute of Himalayan

Environment & Development, Kosi-Katarmal, Almora, for providing necessary facilities to write this article.

## References

- Abbott, D. L. 1984. *The Apple Tree: Physiology and Management*. Growers Association, Kullu Valley, H.P.
- Agarwal, P. K. 2009. *Global Climate Change and Indian Agriculture*. Indian Council of Agricultural Research, New Delhi.
- Ageeta, Y. & T. Kadota. 1992. Predictions of changes of glacier mass balance in the Nepal 36 K. Higuchi Himalaya and Tibetan Plateau: a case study of air temperature increase for three glaciers. *Annals of Glaciology* **16**: 89-94.
- Agnihotri, R. K. & L. M. S. Palni. 2007. On farm conservation of landraces of rice (*Oryza sativa* L.) through cultivation in the Kumaun region of Indian, Central Himalaya. *Journal of Mountain Science* **4**: 354-360.
- Ahmad, F., U. Partap, S. R. Joshi & M. B. Gurung. 2002. Please do not steal our honey. *Bees for Development Journal* **64**: 9.
- Aneja, V. P., S. Businger, Z. Li, C. S. Claiborn & A. Murthy. 1991. Ozone climatology at high elevations in the southern Appalachians. *Journal of Geophysical Research* **96**: 1007-1021.
- Bale, J. S., G. J. Masters, I. D. Hodkinson, C. Awmack, T. M. Bezemer, V. K. Brown, J. Butterfield, A. Buse, J. C. Coulson, J. Farrar, J. E. G. Good, R. Harrington, S. Hartley, T. Hefin Jones, R. L. Lindroth, M. C. Press, I. Symrnioudis, A. D. Watt & J. B. Whittaker. 2002. Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Global Change Biology* **8**: 1-16.
- Bahadur, J. 1998. Himalayan Glaciers. <http://www.mtnforum.org/en/node/6538>.
- Bandyopadhyay, J. & D. Gyawali. 1994. Himalayan water resources: ecological and political aspects of management. *Mountain Research & Development* **14**: 1-24.
- Bhatt, I. D., R. S. Rawal & U. Dhar. 2000. The availability, fruit yield, and harvest of *Myrica esculenta* in Kumaun (west Himalaya), India. *Mountain Research & Development* **20**: 146-153.
- Bhatta, A. 2007. Himachal villagers resist pine monoculture, reclaim forests for fodder. *Down to Earth* **16**: 6.
- Cannone, N., S. Sgorbati & M. Guglielmin. 2007. Unexpected impacts of climate change on alpine vegetation. *Frontiers of Ecology & Environment* **5**: 360-364.
- Chakraborty, S. & S. Datta. 2003. How will plant pathogens adapt to host plant resistance at elevated CO<sub>2</sub> under changing climate? *New Pathologist* **159**: 733-742.
- Dobhal, D. P., J. T. Gergan & R. J. Thayyen. 1999. Recession of dokriani glacier, Garhwal Himalaya - An overview. pp. 30-33. *In: Abstract of Symposium on Snow, Ice and Glaciers. A Himalayan Perspective*. Geological Survey of India, Lucknow.
- Epstein, Y., E. Sohar & Y. Shapiro. 1995. Exceptional heatstroke: A preventable condition. *Israel Journal of Medical Science* **31**: 454-462.
- Eriksson, M., J. Fang, J. & J. Dekens. 2008. 'How does climate affect human health in the Hindu Kush-Himalaya region?' *Regional Health Forum* **12**: 11-15.
- Ghosh, P. & P. P. Dhyani. 2004. Baranaaja: The traditional mixed cropping system of Central Himalaya. *Outlook on Agriculture* **33**: 261-266.
- Guleria, R. P., J. C. Kuniyal, P. S. Rawat, N. L. Sharma & A. K. Thakur. 2010. Aerosols and their radiative forcing over Mohal-Kulu in the Northwestern Indian Himalaya. *In: Proceedings of Aerosols and Clouds: Climate Change Perspective*. IASTA-2010, Bose Institute, Darjeeling, West Bengal, India, Vol. **19**: 198-200.
- Hasnain, S. I. 1999. *Himalayan Glaciers: Hydrology and Hydrochemistry*. Allied Publication Limited, New Delhi.
- Hasnain, S. I. 2002. Himalayan glaciers melt down: impacts on South Asian Rivers. pp. 417-423. *In: H. Lanen, A. J. Van & S. Demuth (eds.) FRIEND 2002-Regional Hydrology: Bridging the Gap Between Research and Practice*. IAHS Publications, Wallingford.
- IPCC. 3rd Assessment Report. 2001a. [http://en.wikipedia.org/wiki/IPCC\\_Third\\_Assessment\\_Report](http://en.wikipedia.org/wiki/IPCC_Third_Assessment_Report).
- IPCC. 2001b. *Climate Change- The Scientific Basis, Summary for Policy Makers and Technical Summary of the Working Group I Report*. Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge.
- IPCC. 2002. *Climate Change and Biodiversity*. Technical Paper V. Intergovernmental Panel on Climate Change. WMO-UNEP.
- Jeelani, Gh. 2008. Aquifer response to regional climate variability in a part of Kashmir Himalaya in India. *Hydrology Journal* **16** : 1625-1633.
- Joshi, S. C. & L. M. S. Palni. 2005. Greater sensitivity of *Hordeum himalayens* Schultz to increasing temperature causes reduction in its cultivated area. *Current Science* **89**: 879-882.
- Joshi, S. C., S. Chandra & L. M. S. Palni. 2007. Differences in photosynthetic characteristics and accumulation of osmoprotectants in saplings of evergreen plants grown inside and outside a glasshouse during the winter season. *Photosynthetica* **45**: 594-600.

- Joshi, V. & G. C. S. Negi. 1995. Analysis of long-term weather data from Garhwal Himalaya. *ENVIS Bulletin on Himalayan Ecology* **3**: 63.
- Kelly, A. E. & M. L. Goulden. 2008. Rapid shifts in plant distribution with recent climate change. *Proceedings of National Academy of Sciences, USA* **105** : 11823-11826.
- Kulkarni, A. V., B. P. Rathore & A. Suja. 2004. Monitoring of glacial mass balance in the Baspa basin using accumulation area ratio method. *Current Science* **86** : 101-106.
- Kumar, R., A. K. Shai, K. Krishna Kumar, S. K. Patwardhan, P. K. Mishra, J. V. Rewadhar, K. Kamal & G. B. Pant. 2006. High resolution climate change scenario for India for the 21st century. *Current Science* **90**: 334-345.
- Kumar, K., S. Joshi & V. Joshi. 2008a. Climate variability, vulnerability, and coping mechanism in Alaknanda catchment, Central Himalaya, India. *AMBIO* **37**: 286-291.
- Kumar, K., R. Dumka, M. S. Miral, G. S. Satyal & M. Pant. 2008b. Estimation of the retreat of gangotri glacier using rapid static and kinematic GPS survey. *Current Science* **94**: 258-261.
- Kuniyal, J. C. 2002. Mountain expeditions: minimizing the impact. *Environmental Impact Assessment Review* **22**: 561-581.
- Kuniyal, J. C., P. S. P. Rao, G. A. Momin, P. D. Safai, S. Tiwari & K. Ali. 2007. Trace gases behaviour in sensitive areas of the northwestern Himalaya: A case study of Kullu-Manali tourist complex, India. *Journal of Radiation & Space Physics* **36**: 197-203.
- Kuniyal, J. C., A. Thakur, H. K. Thakur, S. Sharma, P. Pant, P. S. Rawat & K. K. Moorthy. 2009. Aerosol optical depths at Mohal-Kullu in the northwestern Indian Himalayan high altitude station during ICARB. *Journal of Earth System Science* **118**: 41-48.
- Kuniyal, J. C. 2010. Aerosols climatology over the northwestern Himalayan region. pp. 93-99. *In: Proc. ARFI & ICARB Scientific Progress Report*. ISRO Geosphere Biosphere Programme (IGBP), Space Physics Laboratory, VSSC, Thiruvanthapuram.
- Leemans, R. & B. Eickhout. 2004. Another reason for concern: regional and global impacts on ecosystems for different levels of climate change. *Global Environment Change* **14**: 219-228.
- Liu, X. & B. Chen. 2000. Climate warming in the Tibetan Plateau during recent decades. *International Journal of Climatology* **20**: 1729-1742.
- Maikhuri, R. K., R. L. Semwal, K. S. Rao, S. Nautiyal & K. G. Saxena. 1997. Eroding traditional crop diversity imperils the sustainability of agricultural systems in Central Himalaya. *Current Science* **73**: 777-782.
- Maikhuri, R. K., K. S. Rao & R. L. Semwal. 2001. Changing scenario of Himalayan agroecosystems: loss of agrobiodiversity, an indicator of environmental change in Central Himalaya, India. *The Environmentalist* **21**: 23-39.
- Matny, L. 2000. *Melting of Earth's Ice Cover Reaches New High*. Worldwatch News Brief, 6 March 2000.
- McMichael, A. J., D. H. Campbell-Lendrum, C. F. Corvalan, K. L. Ebi, A. K. Githeko, J. D. Scheraga & A. Woodward (eds). 2003. *Climate Change and Human Health - Risks and Responses*. WHO, Geneva.
- Ming, J., H. Cachier, C. Xiao, D. Qin, S. Kang, S. Hou & J. Xu. 2008. Black carbon record based on a shallow Himalayan ice core and its climatic implications. *Atmospheric Chemistry & Physics* **8**: 1343-1352.
- Musselman, R. C. & W. J. Massman. 1999. Ozone flux to vegetation and its relationship to plant response and ambient air quality standards. *Atmospheric Environment* **33**: 65-73.
- Negi, G. C. S. 1989. *Phenology and Nutrient Dynamics in Tree Leaves of Kumaun Himalayan Forests*. Ph.D. Thesis. Kumaun University, Naini Tal.
- Negi, G. C. S., H. C. Rikhari & S. P. Singh. 1991. Phenological features in relation to growth forms and biomass accumulation in an alpine meadow of the central Himalaya. *Vegetatio* **101**: 161-170.
- Negi, G. C. S. & V. Joshi. 2002. Studies in the western Himalayan micro-watersheds for global change impact assessment and sustainable development. pp. 153-165. *In: K. L. Shrestha (ed.) Global Change and Himalayan Mountains*. Institute for Development and Innovation, Kathmandu, Nepal.
- Negi, G. C. S. & V. Joshi. 2004. Rainfall and spring discharge patterns and relationships in two small catchments in the western Himalayan Mountains, India. *The Environmentalist* **24**: 19-28.
- Negi, G. C. S. & L. M. S. Palni. 2010. Responding to the challenges of climate change: mountain specific issues. pp. 293-307. *In: N. Jeerath, Ram Boojh & G. Singh (eds.) Climate Change, Biodiversity and Ecological Security in the South Asian Region*. MacMillan Publishers India Ltd., New Delhi.
- Nowak, R. S., D. S. Ellsworth & S. D. Smith. 2004. Tansley review: Functional response of plants to elevated atmospheric CO<sub>2</sub> - do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytologist* **162**: 253-280.
- Pant, P., P. Hegde, U. C. Dumka, A. Saha, M. K. Srivastava & R. Sagar. 2006. Aerosol characteristics at a high-altitude location during ISRO-GBP Land Campaign-II. *Current Science* **91**: 1053-1061.
- Pasupalak, S. 2009. Climate change and agriculture in Orissa. *Orissa Review April-May*: 49-52.

- Patterson, D. T., J. K. Westbrook, R. J. V. Joyce, P. D. Lingren & J. Rogasik. 1999. Weeds, insects, and diseases. *Climatic Change* **43**: 711-727.
- Patz, J. A. & W. J. M. Martens. 1996. Climate impacts on vector-borne disease transmission: Global and site-specific. *Journal of Epidemiology* **6**: S145-S148.
- Patz, J. A., D. Campbell-Lendrum, T. Holloway & J. A. Foley. 2005. Impact of regional climate change on human health. *Nature* **438**: 310-317.
- Partap, U. 1999. Conservation of endangered Himalayan honeybee, *Apis cerana* for crop pollination. *Asian Bee Journal* **1**: 44-49.
- Partap, U. & T. Partap. 2003. *Warning Signals from the Apple Valleys of the Hindu Kush - Himalayas: Productivity Concerns and Pollination Problems*. ICIMOD, Kathmandu.
- Pauli, H., M. Gottfried & G. Grabherr. 2003. Effects of climate change on the alpine and nival vegetation of the Alps. *Journal of Mountain Ecology* **7** (Suppl.): 9-12.
- Pendall, E., S. Bridgham, P. J. Hanson, B. Hungate, D. W. Kicklighter, D. W. Johnson, B. E. Law, Y. Luo, J. P. Megonigal, M. Olsrud, M. G. Ryan & S. Wan. 2004. Below-ground process responses to elevated CO<sub>2</sub> and temperature: A discussion of observations, measurement methods, and models. *New Phytologist* **162**: 311-322.
- Porter, J. H., M. L. Parry & T. R. Carter. 1991. The potential effects of climatic change on agricultural insect pests. *Agriculture, Forestry & Meteorology* **57**: 221-240.
- Prasad, P., K. Chauhan, L. S. Kandari, R. K. Maikhuri, A. Purohit, R. P. Bhatt & K. S. Rao. 2002. *Morchella esculenta* (Guchhi): Need for scientific intervention for its cultivation in Central Himalaya. *Current Science* **82**: 1098-1100.
- Raina, V. K. 2009. *Himalayan Glaciers-A State of Art Review of Glacial Studies*. Glacial Retreat and Climate Change. MoEF Discussion paper, GBPIHED & MoEF, Govt. of India.
- Ramakrishna, R. N., C. D. Keeling, H. Hashimoto, W. M. Jolly, S. C. Piper, C. J. Tukder, R. B. Myneni & S. W. Running. 2003. Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *Science* **300**: 1560-1563.
- Ramanathan, V. & G. Carmichael. 2008. Global and regional climate changes due to black carbon. *Nature & Geosciences* **1**: 221-227.
- Ravindranath, N. H., N. V. Joshi, R. Sukumar & A. Saxena. 2008. Impact of climate change on forests in India. *Current Science* **90**: 354-361.
- Reiter, P. 1998. Global warming and vector-borne disease in temperate regions and at high altitude. *Lancet* **351**: 839-840.
- Rosenzweig, C. & D. Hillel. 1995. Potential impacts of climate change on agriculture and food supply. *Consequences* 1, No. 2. (<http://www.gerio.org/CONSEQUENCES/summer95/agriculture.html>: 1-25).
- Rosenzweig, C., A. Iglesias, X. B. Yang, P. R. Epstein & E. Chivian. 2001. Climate change and extreme weather events: Implications for food production, plant diseases, and pests. *Global Change & Human Health* **2**: 90-104.
- Semwal, R. L. & J. P. Mehta. 1996. Ecology of forest fires in Chir Pine forests of Garhwal Himalayas. *Current Science* **70**: 426-427.
- Semwal, R., A. Tewari, G. C. S. Negi, R. Thadani & P. Phartiyal, M. Verma, S. Joshi, G. Godbole & A. Singh (Research Contributors). 2007. *Valuation of Ecosystem Services and Forest Governance: A Scoping Study from Uttarakhand*. Leadership in Environment & Development, LEAD-India, New Delhi.
- Sharma, E., N. Chettri, K. Tse-ring, A. B. Shrestha, F. Jing, P. Mool & M. Eriksson. 2009. *Climate Change Impacts and Vulnerability in the Eastern Himalayas*. International Centre for Integrated Mountain Development, Kathmandu.
- Sharma, S. & H. C. Rikhari. 1997. Forest fire in the central Himalaya: climate and recovery of trees. *International Journal of Biometeorology* **40**: 63-70.
- Shrestha, A. B., C. P. Wake, P. A. Mayewski & J. E. Dibb. 1999. Maximum temperature and trends in the Himalaya and its vicinity: An analysis based on temperature records from Nepal from period 1971-94. *Journal of Climate* **12**: 2775-2787.
- Shukla, S. P. & M. A. Siddiqui. 1999. Recession of the snout front of Milam Glacier, Goriganga valley, District Pithoragah, Uttar Pradesh. pp. 27-29. *In: Abstract of Symposium on Snow, Ice and Glaciers. A Himalayan Perspective*. Geological Survey of India, Lucknow.
- Singh, S. P., V. Singh & M. Skutsch. 2010. Rapid warming in the Himalayas: Ecosystem responses and development options. *Climate & Development* **2**: 1-13.
- Sinha, S. 2007. Impact of climate change in the highland agroecological region of India, Sahara Time Magazine, (<http://www.saharatime.com/Newsdetail.aspx?newsid=2659>).
- Thakur, V. C., N. S. Viridi, J. T. Gergan, R. K. Mazari, R. K. Chaujar, S. K. Bartarya & G. Philip. 1991. *Report on Gaumukh: The Snout of the Gangotri Glacier*. Unpublished Report submitted to DST, New Delhi.
- Thorntwaite, C. W. 1948. An approach towards a rational classification of climate. *Geographical Review* **38**: 55-94.
- Tong, S. L. & L. V. Ying. 2000. Global change and epi-

- demic disease. *Journal of Disease Control* **4**: 17-19.
- Uprety, D. C. & V. R. Reddy (eds.). 2008. *Rising Atmospheric Carbon Dioxide and Crops*. Indian Council of Agricultural Research, New Delhi.
- USEPA. 1996. *Air Quality Criteria for Ozone and Related Photochemical Oxidants*. U. S. EPA, Research Triangle Park, NC EPA Report No. EPA/600/P-93/00aF, Vol.II.
- Valdiya, K. S. & S. K. Bartarya. 1989a. Diminishing discharge of mountain springs in a part of Kumaun Himalaya. *Current Science* **58**: 417-424.
- Valdiya, K. S. & S. K. Bartarya. 1989b. Problem of mass movements in a part of Kumaun Himalaya. *Current Science* **58** : 486-491.
- Vedwan, N. & R. E. Rhoades. 2001. Climate change in the western Himalayas of India: a study of local perception and response. *Climate Research* **19**: 109-117.
- Vishvakarma, S. C. R., J. C. Kuniyal & K. S. Rao. 2003. Climate change and its impact on apple cropping in Kullu Valley, North-West Himalaya, India. In: *7th International Symposium on Temperate Zone Fruits in the Tropics and Subtropics*. 14-18 October, Nauni-Solan (H.P.).
- Vohra, C. P. 1981. Note on recent glaciological expedition in Himachal Pradesh. *Geological Survey of India Special Publication* **6**: 26-29.
- Walter, H. 1985. *Vegetation Systems of the Earth and Ecological Systems of the Geo-Biosphere*. Springer-Verlag, Berlin.
- Watson, R. T., M. C. Zinyowera & R. H. Moss (eds.). 1997. Climate change 1995; impacts, adaptations and mitigation of climate change: scientific-technical analysis. *Contribution of Working Group II to the Second Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, Cambridge University Press.
- Whittaker, R. H. 1975. *Communities and Ecosystems*. Macmillan, New York.
- World Climate News. 2006. *Homing in on Rising Sea Levels*. WMO, Geneva, Switzerland, No. 29, June **29**: 1-12.
- Woodruff, R. E. & C. S. Guest. 2002. Predicting Ross River virus epidemics from regional weather data. *Epidemiology* **13**: 384-393.

(Received on 23.09.2010 and accepted after revisions, on 04.10.2011)